


Date & Time of Appointment _____

Please mark the figure with an "x" everywhere you have pain



Time of
Injection

Worst ever
SI joint pain

/10

SI joint pain
today

/10

For each time period, please shade the circle that corresponds
to your SI joint pain level and include any remarks:

Worst Pain Imaginable	Before injection	After injection					
		30 min.	1 hr.	90 min.	2 hr.	3 hr.	4 hr.
10	☐	☐	☐	☐	☐	☐	☐
9	☐	☐	☐	☐	☐	☐	☐
8	☐	☐	☐	☐	☐	☐	☐
7	☐	☐	☐	☐	☐	☐	☐
6	☐	☐	☐	☐	☐	☐	☐
5	☐	☐	☐	☐	☐	☐	☐
4	☐	☐	☐	☐	☐	☐	☐
3	☐	☐	☐	☐	☐	☐	☐
2	☐	☐	☐	☐	☐	☐	☐
1	☐	☐	☐	☐	☐	☐	☐
No Pain 0	☐	☐	☐	☐	☐	☐	☐

Remarks _____

Si-Logix

If sacroiliac joint pain limits your life, the
SacroFuse procedure can help you move with
comfort again.

Ask your healthcare provider if this minimally
invasive option is right for you.

For more information, talk with your physician
or visit your clinic's website.

SacroLoX


Take Back Your
Movement

Si-Logix | **SacroLoX**
A Division of Evologics

www.si-logix.com



1. Understanding SI-Joint Pain



1 Your sacroiliac

(SI) joints connect the base of your spine (sacrum) to your pelvis. When these joints become irritated or unstable, it can cause pain that affects daily life.

2 What SI Joint Pain Feels Like

- Aching or sharp pain in the lower back, buttocks, or hips
- Discomfort that radiates to the groin or legs
- Pain that worsens with standing or walking
- Difficulty sitting for long periods or sleeping comfortably


Common Experiences & Symptoms

People with SI joint dysfunction can experience pain, weakness, and limited mobility.

Common Symptoms:

- Lower back pain (below L5)
- Pelvis or buttock pain
- Hip or groin pain
- Leg weakness or instability
- Numbness or tingling in the legs
- Pain when moving from sitting to standing
- Trouble sleeping due to pain
- Difficulty sitting for long periods

Your doctor can confirm SI joint dysfunction with a specific examination.



About the SacroFuse Procedure

• Procedure Highlights

- Minimally invasive approach
- Two cylindrical allograft implants inserted through a bone access needle
- Provides immediate joint stabilization and promotes fusion
- Usually performed under sedation in less than an hour

• After the Procedure

- Most patients walk the same day
- Return to light or non-strenuous work in about two weeks
- Follow your healthcare provider's postoperative instructions carefully

